

# Pathway for the Delirium Module of the AO Trauma Orthogeriatrics App


**Katrin Singler<sup>1,2</sup>, Markus Gosch<sup>2</sup>, Tobias Roth<sup>3</sup>, Sacha Beck<sup>4</sup>**

<sup>1</sup>Institute for Biomedicine of Aging, Friedrich-Alexander-University Erlangen-Nürnberg, Germany  
<sup>2</sup>Klinikum Nürnberg, Paracelsus Private Medical University, Germany  
<sup>3</sup>Medical University of Innsbruck, Austria  
<sup>4</sup>Stadtspital Waid, Zürich, Switzerland



**Responsible use of phones and tablets:** These devices are a potential source of infection, distraction, and patient dissatisfaction. Consult your hospital regulations before using devices in the clinical setting.

## An approach to delirium in older adults with a fracture (excluding substance-withdrawal delirium)



**References**

**Inouye SK, VanDyck CH, Alessi CA, et al.** Clarifying confusion: The Confusion Assessment Method. A new method for detecting delirium. *Ann Intern Med.* 1990; 113:941-948.  
**Vochteloo AJ, Moerman S, van der Burg BL, et al.** Delirium risk screening and haloperidol prophylaxis program in hip fracture patients is a helpful tool in identifying high-risk patients, but does not reduce the incidence of delirium. *BMC Geriatr.* 2011 Aug 11; 11:39.

**Notes** (for local protocols, dosages, etc)